

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 43/14, 69/618, 69/52, A61K 31/215, 31/075	A1	(11) International Publication Number: WO 95/16659 (43) International Publication Date: 22 June 1995 (22.06.95)
(21) International Application Number: PCT/US94/12130 (22) International Filing Date: 20 October 1994 (20.10.94) (30) Priority Data: 08/167,424 15 December 1993 (15.12.93) US (71) Applicant: AVON PRODUCTS, INC. [US/US]; Division Street, Suffern, NY 10901 (US). (72) Inventors: DUFFY, John, A.; 17 Yorkshire Avenue, West Milford, NJ 07480 (US). TEAL, Janice, J.; 70 Bon Aire Circle, Suffern, NY 10910 (US). GARRISON, Mark, S.; 12 Tomahawk Drive, White Plains, NY 10603 (US). SERBAN, George, P.; 239 Limestone Road, Ridgefield, Conn. 06877 (US). (74) Agent: TODD, William, G.; Hopgood, Calimafde, Kalil & Judlowe, 60 East 42nd Street, New York, NY 10165 (US).	(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LT, LV, MD, MG, MN, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report.</i>	
(54) Title: NOVEL RETINOID CONJUGATE COMPOUNDS AND METHODS FOR TREATING SKIN AGING		
(57) Abstract		
Skin aging and related conditions can be effectively treated with a conjugate of a retinoid and a bioactive organic acid preferably selected from among alpha-hydroxy acids, beta-hydroxy acids, and keto-acids; preferred conjugates include retinyl glycolic ether and retinyl glycolate (as either the ester or reverse ester).		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

NOVEL RETINOID CONJUGATE COMPOUNDS
AND METHODS FOR TREATING SKIN AGING

Background of the Invention

1. The Field of the Invention

5 The present invention relates to conjugates of retinoids with organic acids and to the use of such compounds in the treatment of skin aging.

2. The State of the Art


10 Skin aging is accompanied by a number of morpho-physiological changes which are described in the literature. See, e.g., B.A. Gilchrest, J. Cutaneous Aging Cosmet. Dermatol., 1:1-3 (1988), and Arch. Dermatol., 115:1343-1346 (1979). At the physiological level, skin aging is accompanied by a decrease in at least one
15 critical skin function, such as epidermal turnover, healing, clearance of chemicals from the dermis, water regulation, sensory perception, mechanical protection, immunocompetence, vascular reactivity, blood flow, sweat gland function, turgor, sebum production or Vitamin D
20 synthesis.

 At the morphological level, a decline in critical physiological functions is associated with a loss in the normal pattern of corneocyte desquamation, epidermal thinning and/or the presence of atypical epidermal cells.
25 Other characteristics include a reduction in the height of rete ridges along with a corresponding effacement of dermal papillae, a general weakening of the dermo-epidermal junction, excessive accumulation of elastin and a loss of collagen and other ground substances, a decrease
30 in dermal volume (tugor), pigmentary disorders, attrition

of capillary vessels, chronic inflammation and the occurrence of benign and/or malignant tumors. Some of the clinical manifestations of these morphological changes which are relevant to cosmetic appearance and dermal health include a rough, flaky and/or dry skin surface, itching, excessive wrinkling, sagging, loss of elasticity, 5 sallow color, mottled pigmentation, thinning hair, nail brittleness and, in some cases, skin growths.

The etiological bases underlying skin aging are only partially understood. But, it is recognized that the effect of ultraviolet light (UVL), especially from the sun, constitutes a significant factor in the acceleration of skin aging. Several studies have shown that skin which has been exposed to sunlight undergoes aging sooner than unexposed areas in the same individual. By comparing the histological and functional characteristics of exposed and unexposed skin, investigators have observed differences between chronological and accelerated premature aging. See, e.g., W. Montagna, J. Invest. Dermatol., 73:47-53 (1979); B.A. Gilchrest, J. Invest. Dermatol., 80:81s-85s (1983); and B.A. Gilchrest, J. Invest. Dermatol., 73:59-66 (1979).

The UVL component of sunlight is responsible for triggering molecular changes that lead to damaged biological tissues, including the skin. It is postulated that one mechanism for sun-induced aging results from the generation of free radicals in the skin. One proposed pathway to free radical generation, the so-called Haber-Weiss reaction (shown in the scheme below) is a chemical pathway which generates the most reactive free radical species, the hydroxy radical.





wherein $\frac{1}{2}O_2$ is singlet oxygen or the superoxide ion, $HO_2\cdot$ is the hydroperoxyl radical, H_2O_2 is hydrogen peroxide, $HO\cdot$ is the hydroxy radical and H_2O is water. In this reaction, electrons (e^-) are gained through the oxidation of metals such as iron, copper and zinc, which are typically present in the skin. The hydrogen atoms are gained through the destructive oxidation of biological molecules essential to the normal function of cells, tissues and organs, such as lipids, proteins, DNA, RNA, and enzymatic cofactors. There are numerous studies showing how the oxidation of these molecules is detrimental to the skin, in both the long term and short term.

The retinoids are biologically active compounds involved in essential functions such as vision, embryonic development and the growth and maintenance of normal skin. The art teaches that topical application of skin preparations containing retinoids provides a significant improvement in clinical appearance. See, for example, U.S. Patent Nos. 4,603,146 and 4,877,805. Additionally, topical retinoid application provides improvement with respect to several histological parameters, such as thickening of the epidermis including the stratum granulosum, an increase in the height of rete ridges and the number of dermal papillae, a gradual displacement of age-related deposition of dermal elastin by collagen and peptidoaminoglycans, normalization of melanocyte function and an increase in the number of dermal fibroblasts. See, e.g., A.S. Zelickson, J. Cutaneous Aging Cosmet. Dermatol., 1:41-47 (1988); J.S. Weiss, JAMA, 259:527-532 (1988); J. Bhawan, Arch. Dermatol., 127:666-672 (1991); and L.H. Kligman, Connect. Tissue Res., 12:139-150 (1984). U.S. Patent No. 4,603,146 also describes the use of Vitamin A acid for retarding skin aging, and U.S. Patent No. 4,877,805 describes the use of retinoids generally for the same purpose (the disclosures of which are both incorporated herein by reference).

The skin is a target for and requires retinoids. This conclusion stems from the fact that Vitamin A deficiencies lead to skin lesions. Additionally, skin cells, such as fibroblasts and keratinocytes, contain high-affinity cytosolic receptors for retinoids, thus further demonstrating the target nature of these skin cells.

It is well-known that therapeutic doses of topically applied retinoids frequently cause skin irritations which interfere with treatment. To solve this problem, the art has resorted to conjugating retinoids with glucuronic acid. For example, U.S. Patent No. 4,855,463 discloses water-soluble glucuronic acid derivatives of Vitamin A (formula I) for improved metabolic uptake; while S. Chen, J. Invest. Dermatol., Abstract, 98:560 (1992) describes acetaminophen (formula II). In both cases, conjugation was achieved by attaching the other molecule to carbon-15 of the retinol structure.



U.S. Patent No. 4,216,224 describes retinyl esters of hydroxy acids, hydroxy amides or hydroxy acid esters for use in treating psoriasis; and U.S. Patent No. 5,124,356 describes ester and amide conjugates of trans-retinoids, although the molecule conjugated with the retinoid is not described as providing any separate therapeutic benefit. EP-A2-0 391 033 further describes the preparation of

retinal derivatives, especially acetals and hemiacetals, useful in treating a variety of skin conditions.

JP-A 63-66160/1980 describes the preparation of a retinoyl ester of 1-ascorbic acid which is taught to be useful for increasing the activity of blood vessel endotheliocyte plasminogen activators, for increasing the hydrophilic properties of retinol and retinoic acid to reduce instability of the composition due to oxidation. In vitro data on endotheliocyte cultures showed increased activity of plasminogen activator, but no methods or compositions for administration are disclosed.

JP-A 4-210686/1992 describes a method for the manufacture of tocopheryl retinoate. This compound is useful for treating skin ulcers (e.g., JP-A 61-207332) and for preventing the roughening of skin (e.g., JP-A 51-73137). U.S. Patent No. 5,182,396 discloses that 1-hydroxy Vitamin D esters with Vitamin A acid are useful for treating cutaneous ulcers and tumors.

Carboxylic acids are also useful in the treatment of aging skin, especially alpha- and beta-hydroxy acids and keto-acids. These are generally referred to as AHAs (Alpha-Hydroxy Acids). See, e.g., U.S. Patent Nos. 3,920,835, 4,045,559, 4,053,630 and 4,363,815, the disclosures of which are incorporated herein by reference. Additionally, U.S. Patent No. 4,194,007 describes α -hydroxyretinoic acid and α -ketoretinoic acid derivatives of retinoic acid.

These AHAs improve the clinical appearance and mechanical properties of aging skin. A normalization in the pattern of epidermal keratinization and a reduction in the cohesive forces acting between keratinocytes and corneocytes have also been reported after treatment with AHAs (E.J. Van Scott, J. Acad. Dermatol., 11:867-879 (1984)). Other investigators have observed such benefits as wrinkle reduction after topical application of AHAs. They postulate that AHAs stimulate collagen and glycosaminoglycan synthesis by cultured fibroblasts (F.D.

Dial, Cosmetic Dermatology, 5:32-34 (1990)). AHAs also increase the production of collagen and peptidoaminoglycans (R.M. Lavker, J. Am. Acad. Dermatol., 26:535-544 (1992)), thereby further improving the general topographical appearance and viscoelastic behavior of skin.

The action by which AHAs improve the clinical appearance and mechanical properties of skin is not well-understood. One feature of aging skin is hyperkeratinization, a condition in which the corneocytes adhere in excess, thereby accounting for a thickened stratum corneum and a dry appearance. The forces responsible for the adhesion of corneocytes are non-covalent in nature and evidence exists that both ionic and hydrophobic interactions are involved.

In the case of ionic interactions, calcium ions are believed to bridge adjacent corneocytes by forming complexation bonds between cholesteryl sulfate residues that are firmly anchored to the corneocyte cell wall (L.J. Shapiro, Lancet, 1:70-72 (1978)). Since ionic interactions are Ph dependent, AHAs may decrease corneocyte cohesion by temporarily opening up calcium bridges. However, an explanation of activity based on Ph alone is not satisfactory. Earlier work aimed at studying the relationship between the chemical structure of these agents and their ability to influence the viscoelastic behavior of the stratum corneum clearly indicates that, among the series of AHAs studied, those having hydroxy groups in the alpha position consistently lowered this parameter (M. Takahashi and Y. Machida, J. Soc. Cosmet. Chem., 36:177-187 (1985)). The lower the viscoelastic modulus, the less force is required to cause deformation (i.e., the softer the stratum corneum).

AHAs are known to produce a long-lasting decrease in the elastic modulus of skin. This means softening and, with regular use, a normalization in the pattern of corneocyte desquamation. Recent work shows that these

beneficial effects are optimized after the AHAs are conjugated with linear aliphatic chains having eight to ten carbons (e.g., D.B. Hagan et al., "A study of the structure-activity relationships present in skin active agents," Int'l J. Cosmetic Sci., 15, 163-173 (1993), the disclosure of which is incorporated herein by reference).

It will be appreciated that a range of compounds have been considered for the treatment of skin aging. While there are theoretical, if not observed, benefits derived from these compounds, problems relating to stability; toxicity and dermal penetration still exist. These problems limit the availability of useful therapies for skin aging.

Summary of the Invention

It is among the objects of this invention to provide a new class of compounds useful in the treatment of aging skin and to provide methods for the use of such compounds in the treatment of aged and aging skin. Particular benefits achieved by the present invention include increased stability of the active compound, improved dermal penetration and a higher therapeutic index (i.e., selectively, typically defined as TD50/ED50 or LD50/ED50).

The present invention specifically provides a class of multi-functional conjugated retinoids, bimers of a retinoid and other bioactive ingredients selected from the group consisting of chelating agents, anti-oxidants and low molecular weight organic acids. When used for treating aged skin, these novel conjugates provide a higher therapeutic index than any of the classes of agents discussed in the Background section of this disclosure, whether applied individually or in combination.


The present invention also provides a pharmaceutical composition which comprises an amount of the novel conjugate effective to provide a benefit to a morphological and/or clinical aspect of aging skin and a compatible carrier therefor; these aspects are described

above in the Background of section. In a preferred embodiment, the carrier is cosmetically acceptable, if not dermatologically acceptable.

The invention also provides a method for treating aging skin of a patient having such aged skin, which comprises providing the novel conjugate of this invention as a composition comprising a mixture of the conjugate and a cosmetically acceptable carrier therefor, and applying said composition to the patient such that the amount of conjugate applied is effect to benefit a morphological and/or clinical aspect of the patient's aging skin.

Detailed Description of Specific Embodiments

The present invention is directed to novel conjugates of retinoids. As used herein, the term "retinoid" includes Vitamin A (retinol) and derivatives such as retinoic acid (e.g., tretinoin which is available as Retin-ATM from Johnson & Johnson Co., New Brunswick, New Jersey, USA), retinal (Vitamin A aldehyde), 3,4-didehydroretinol (Vitamin A₂); and cosmetically acceptable derivatives such as Vitamin A acetate and Vitamin A palmitate, and other esters or reverse esters (or salts thereof), ethers, aldehydes, alcohols, and the like. Retinol has the structure of formula III which is shown below.



The retinoid is conjugated with a bioactive agent selected from organic acids, especially those which are cosmetically acceptable, and particularly those which have a beneficial effect on the appearance and/or properties of aging skin. It is preferred that the retinoid be conjugated with a low-molecular weight organic acid (i.e.,

those having not more than about 10 carbon atoms). It is preferred that the linkage between the retinoid and the organic acid be through an ester, reverse ester, ether or amide bond, and most preferred that the linkage be an ether bond. It is believed that these types of bonds can be cleaved by non-specific, ubiquitous hydrolytic enzymes present in large amounts in the viable dermal tissues, and to a lesser extent in the stratum corneum. While not desirous of being constrained to a particular theory, after delivery to the dermal tissues, these novel conjugates are believed to hydrolyze in vivo and yield an active retinoid and active AHA agent. Both compounds provide anti-aging effects when made available to the skin.

The novel conjugates of this invention can be made using organic acids or their derivatives, such as aldehydes, ketones, alcohols, esters, reverse esters, anhydrides, acyl halides, and salts thereof, and preferably those which are bioactive (although, in some situations, it may be desirable to provide an organic acid which is relatively inert). The organic acid preferably has from 2 to 24 carbon atoms, more preferably from 4 to 18 carbon atoms, most preferably from 3 to 12 carbon atoms, and especially including 6 to 10 carbon atoms. Particular α -hydroxy organic acids suitable for use with this invention include, as examples, glycolic, lactic, citric, tartaric, mandelic, benzoic, and malic acids. Particular oxoacids suitable for use with this invention include, for example, glyoxylic and pyruvic acids. Other carboxylic acids, such as succinic, fumaric, oxalic, and salicylic acids, are also suitable for use with this invention. Other suitable organic acids include those which are normal cellular constituents. More preferably, the organic acid is selected from among alpha-hydroxy acids, beta-hydroxy acids, and keto-acids (generally known as AHAs). AHAs generally contain from two to five carbon

atoms in their aliphatic residue and are highly ionized; that is, they are strongly acidic.

Derivatives of organic acids which are suitable in this invention include substitutions on the acid, and especially substitutions at the 2-position (e.g. the α -position of an AHA). Exemplary derivatives include 2-thio derivatives (e.g., wherein the hydroxyl group of the AHA is substituted with a thio group), thioalkyl and thioalkenyl derivatives (e.g., those substituted at the 2-position of a carboxylic acid, and especially of a hydroxycarboxylic acid), keto derivatives, methoxy derivatives, and halogen derivatives (including F, Cl, Br, and I).

The organic acid may generally be saturated, unsaturated, or polyunsaturated, with double bonds independently in cis- or trans- configurations. Further, the organic acid can be straight- or branched-chain, substituted or unsubstituted, acyclic, cyclic, or heterocyclic, and including aromatic compounds. Also generally suitable are organic acids which are liquid at ambient pressure and temperature (i.e., about 25° C. and 1 atm.).

After topical application of an AHA, skin surface pH is temporarily lowered from a normal level of about 5 to a more acidic level of about 3. The localized acidity can result in stinging, particularly on the face where microfissures are more common than on other parts of the body. Because charged molecules penetrate the cutaneous barrier very slowly, it may require several hours for the increased surface acidity to be neutralized, thereby ending the stinging and discomfort to the patient.

Conjugation of a short chain carboxylic acid with a retinoid, especially where the acid is conjugated via the carboxylate moiety, largely quenches the ability of the acidic residue to become deprotonated. Accordingly, the novel conjugates of this invention provide improved treatment by alleviating acidification of the skin and the

accompanying stinging and other discomfort, even when applied to the facial areas.

Charged molecules, such as AHAs, are known to be poor skin penetrators because, ordinarily, they cannot partition and diffuse through the lipid-rich lamellae present in the intercorneocyte spaces. Water-soluble AHAs have low rates of percutaneous absorption for this reason. By conjugating these acids with the larger, more lipophilic retinoid, they tend to pass through the intact domains of the stratum corneum rather than partition from the formulation into the microfissures.

The synthesis of the novel conjugates of this invention may require conjugating the lipophilic retinoid with a lipophobic molecule such as any of the AHAs. Conjugates having ester or even ether linkages may be hydrolyzed by the presence of water, thereby making synthesis more difficult. Accordingly, it can be beneficial if not desirable to protect the hydroxyl and other active moieties on the AHA conjugated with the retinoid. This precaution facilitates formation of the conjugate bond and/or facilitates solubilizing both of the reactants (retinoid and AHA) in a common medium. For example, whereas glycolic acid (hydroxy ethanoic acid) and acetic acid (ethanoic acid) are both soluble in water, the bromoacetic acid methyl ester used in the synthesis exemplified above is insoluble in water, but soluble in alcohol and ether suitable as the reaction medium.

In various synthesis of conjugates within the scope of this invention, it may be necessary to protect one or more reactive groups, such as hydroxyl groups. Such methods of protection for particular substituents are known to those skilled in the arts of organic synthesis.

In practicing the present invention, it should be appreciated that conjugate formation may require a precursor or derivative of the agent required for the synthesis. The skilled artisan can readily choose a

synthesis scheme and a suitable precursor or derivative for the production of the desired conjugate.

In preparing a composition for use in treating skin aging, one or more conjugates (which are chemically compatible) are provided in a cosmetically or pharmaceutically acceptable vehicle. The amount of the conjugate will preferably range between 0.001% and 4% by weight (unless otherwise noted, all fractional amounts are expressed in weight percent), more preferably between about 1.01% and 1.0%, and most preferably between about 0.05% and 0.25% of the composition. The composition should be applied between one and three times each day to an affected area.

A cosmetically acceptable organic acid can optionally be present independently in the composition in an amount, preferably a bioactively effective amount, of 0.1% to 5.0%; the addition of an AHA is a preferred embodiment. When one or more such organic acids are present in the composition, the pH of the composition may need to be raised to more neutral levels by the addition of an alkaline material (e.g., sodium, potassium, or ammonium hydroxide).

Suitable vehicles or carriers for storage and/or delivery of the novel conjugates of this invention may be provided in liquid, ointment, salve, spray, poultice or other forms, and will preferably have a lipophilic character. Suitable carriers include ethanol, petrolatum, triglycerides, various esters, fatty alcohols, and alkylene glycols of which polypropylene glycol and polyethylene glycol are most preferred; if desired, compatible combinations of these vehicles are also suitable.


Carrier systems for topical delivery include lotions, ointments, salves, creams, gels, foams, sprays (both mists and aerosols), patches, masks and the like. The vehicles are present as needed for the desired delivery system. Additional components may be added according to

conventional practice. For example, the final composition may contain various colorants, fragrances, thickeners (such as xanthan gum), preservatives, humectants, surfactants, dispersants and the like, including typical botanical extracts such as those derived from Witch Hazel, chamomile, and the like (e.g., those having an astringent, antiseptic, or other desired effect). The composition may likewise include a penetration enhancer such as DMSO (dimethyl sulfoxide). It may also include one or more additional active ingredients (such as an antibiotic, anesthetic or growth factor).

The invention will be further illustrated by means of the following example, which is not meant to limit this invention to the particular materials, conditions and products described.

EXAMPLE 1

Preparation of Retinoid/Glycolic Acid Ether Conjugate



In a dark room under dim red light, retinol acetate (120 g., 366 mmol, obtained from BASF Corp., Parsippany, NJ) was dissolved in 1.5 L of methanol under an argon atmosphere. Solid potassium carbonate (151.5 g., 1098 mmol) was added with stirring. After seven hours of stirring, the reaction mixture was filtered through a coarse fritted filter to remove solids. Methanol was then removed by evaporation. Diethyl ether was added and the mixture was washed first with distilled water (although tap water could be used), and then washed with brine (i.e., a saturated sodium chloride solution). The washed ether layer was dried over MgSO_4 and Na_2SO_4 , filtered and

distilled under reduced pressure to produce retinol (101 g., 96% yield). Residual oil was crystallized from hexane using a seed crystal of authentic retinol. This general synthesis of retinol from retinol acetate has been described by J.J. Plattner, J. Am. Chem. Soc., 94, 8613 (1972).

Under dim red light, sodium hydride (60% dispersion in mineral oil, 18.4 g., 459 mmol) was placed in an r.b. flask under argon. Hexane (freshly distilled from CaH_2) was used to wash the sodium hydride three times to remove residual mineral oil. Tetrahydrofuran (THF, 350 ml) was added to the sodium hydride with stirring to obtain a suspension. The retinol (101 g.) produced in the preceding step, plus 19 g. of previously obtained retinol crystals (total 120 g., 418 mmol) were dissolved in 300 ml THF. The THF solution of retinol was slowly added to the sodium hydride suspension while the suspension was stirred and maintained in an ice bath. Evolution of hydrogen gas was noted. After this addition, the ice bath was removed and stirring was continued for approximately 30 minutes. A dropping funnel was charged with methyl bromoacetate (97%, 41.76 ml, 430 mmol) and 100 ml THF. The methyl bromoacetate solution was added drop-wise at such a rate that the reaction temperature was maintained at not greater than approximately 30° C. This procedure required approximately two hours and the mixture was stirred for an additional two hours. The product was retinyl acetate ether which resulted from alkylation of the previously formed retinol. The alkylation of retinol with methyl bromide has been described by B.A. Stoochnoff, Tetrahedron Lett., Vol. 21 (1973) and C.A. Brown, Synthesis, Vol. 434 (1974).

Under dim red light, any residual sodium hydride remaining after the alkylation step was quenched by the addition of a few milliliters of methanol. Water (400 ml) was carefully added to the reaction mixture. Lithium hydroxide (30 g., 1254 mmol) was then added. After

15

15 minutes, the reaction was homogeneous. After stirring for two additional hours, the reaction flask was placed on a rotary evaporator to partially remove the THF. The basic aqueous phase was extracted three times with ether to remove the non-base soluble materials, such as unreacted retinol. The aqueous layer was then acidified to a pH of 3 with 4 N HCL at 0° C. and extracted four times with ether to give the crude saponified product (79 g., 220 mmol; 53% yield). Analysis by NMR, UV-IVS and mass spectroscopy confirmed that the product was retinol glycolic ether (retinyl glycolyl ether).

Examples 2-6

The retinyl glycolic ether produced in claim 1 is made into a variety of cosmetically acceptable formulations, including a cream, a gel, a lotion, a toner, and an ointment as shown below. All amounts given are weight fractions unless otherwise noted.

Cream

	Cetyl alcohol	2.00
20	Glyceryl monostearate	2.50
	White Petrolatum	5.00
	Octyl Palmitate	8.00
	Steareth-2	1.00
	PEG 40 Stearate	1.50
25	CARBOMER 940	0.75
	Triethanolamine (99%)	0.75
	Retinyl Glycolyl Ether	0.50
	Water	q.s. to 100%

Gel

30	Propylene Glycol	5.00
	Glycerin	3.00
	Alcohol	10.00
	CARBOMER 940	0.50
	Triethanolamine	0.50

16

Retinyl Glycolyl Ether	2.00
Water	q.s. to 100%

Lotion

	Xantham Gum	0.25
5	Hydroxyethyl cellulose	0.40
	Propylene Glycol	5.00
	Caprylic/Capric Triglyceride	8.00
	Steareth-2	1.50
	PEG 40 Stearate	2.00
10	Retinyl Glycolyl Ether	0.20
	Glycolic Acid	2.00
	Ammonium Hydroxide	q.s. to $3 \leq \text{pH} \leq 4$
	Water	q.s. to 100%

Toner

15	SD alcohol	35.00
	Glycerin	3.00
	Disodium EDTA	0.20
	Botanic Extracts	1.00
	Retinyl Glycolyl Ether	0.01
20	Lactic Acid	1.00
	Ammonium Hydroxide	q.s. to $3 \leq \text{pH} \leq 4$
	Water	q.s. to 100%

Ointment

	White Petroleum	97.6
25	Retinyl Glycolyl Ether	2.40

Various modifications and alterations to the present invention may be appreciated based on a review of this disclosure. These changes and additions are intended to be within the scope and spirit of this invention as defined by the following claims.

WHAT IS CLAIMED IS:

1. A method for the treatment of aged and aging skin, which comprises topical application of an effective amount of a conjugate of a retinoid with a bioactive compound or derivative thereof selected from the group consisting of organic acids having from 2 to 24 carbon atoms.
2. The method as defined by claim 1, wherein the organic acid is selected from the group consisting of α -hydroxy acids, β -hydroxy acids, and keto-acids.
3. The method as defined by claim 1, wherein the organic acid is an α -hydroxy acid.
4. The method as defined by claim 1, wherein the α -hydroxy acid is an acid selected from the group consisting of glycolic, lactic, citric, succinic, fumaric, oxalic and malic acids and derivatives and compatible mixtures thereof.
5. The method as defined by claim 1, wherein the conjugate is an ester, reverse ester, or ether between a retinoid or derivative thereof and a compound selected from the group consisting of glycolic acid, lactic acid, citric acid, succinic acid, fumaric acid, oxalic acid, and malic acid, and derivatives thereof.
6. The method as defined by claim 1, wherein said conjugate is applied in the form of a composition comprising 0.05% to 5% of said conjugate and an acceptable topical carrier therefor.
7. The method as defined by claim 6, wherein said composition comprises 0.3% to 0.6% of said conjugate.

8. A compound useful for the treatment of aged and aging skin which comprises a retinoid conjugated through an ester, reverse ester, ether, or amide linkage with a bioactive compound selected from the group consisting of organic acids and derivatives thereof having from 2 to 24 carbon atoms, to the exclusion of the retinol ester of glucuronic acid.

9. The compound defined by claim 8, wherein the conjugate is an ester of a retinoid and an organic acid.

10. The compound defined by claim 8, wherein the conjugate is an ether of a retinoid and an organic acid.

11. The compound defined by claim 9, wherein the conjugate is a reverse ester of a retinoid and an organic acid.

12. The compound defined by claim 9, wherein the conjugate is an amide of a retinoid and an organic acid.

13. The compound defined by claim 10, wherein the conjugate is a retinyl ester of an α -hydroxy acid selected from the group consisting of glycolic, lactic, citric, succinic, fumaric, oxalic and malic acids and derivatives and compatible mixtures thereof.

14. The compound defined by claim 13, wherein the conjugate is retinyl glycolate.

15. A composition useful for the treatment of aged and aging skin which comprises an effective amount of a compound as defined by claim 9 and a pharmaceutically and cosmetically acceptable carrier therefor.

16. The composition defined by claim 15, wherein the compound is selected from the group consisting of

retinyl glycolate,
retinyl glycolyl ether,
retinyl lactate,
retinyl lactyl ether,
5 retinyl diphenyl glycolate,
retinyl diphenyl glycolyl ether,
retinyl mandelate,
retinyl mandelyl ether,
retinyl citrate,
10 retinyl citryl ether,
retinyl tartrate,
retinyl tartryl ether,
retinyl salicylate,
retinyl salicyl ether,
15 retinyl malate,
retinyl malyl ether,
retinyl succinate,
retinyl fumarate,
retinyl oxalate,
20 retinyl pyruvate, and
compatible mixtures thereof.

17. The composition defined by claim 16, wherein the compound is selected from the group consisting of
retinyl glycolate,
retinyl glycolyl ether,
5 retinyl lactate,
retinyl lactyl ether,
retinyl diphenyl glycolate,
retinyl diphenyl glycolyl ether,
retinyl mandelate,
10 retinyl mandelyl ether,
retinyl salicylate,
retinyl salicyl ether, and
compatible mixtures thereof.

18. The composition as defined by claim 17, wherein the compound is retinyl glycolate or retinyl glycolyl ether.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/12130

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :C07C 43/14, 69/618, 69/52; A61K 31/215, 31/075

US CL :514/847; 568/626

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/847; 568/626

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS Online

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 5,153,230 (JAFFERY) 06 October 1992, entire document, especially the abstract, claims 1-7, and column 2, lines 14-48.	1-18

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be of particular relevance	* X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* &*	document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means		
* P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

28 NOVEMBER 1994

Date of mailing of the international search report

13 JAN 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

MICHAEL B. HYDORN

Facsimile No. (703) 305-3230

Telephone No. (703) 305-1235